




New Cases 


Age-adjusted rate per 100,000 people

Top 5		
1	Utah	26.3
2	New Mexico	38.8
3	Colorado	41.7
4	California	41.7
5	Wyoming	42.4
Bottom 5		
47	Tennessee	74.8
48	Mississippi	75.2
49	Arkansas	77.7
50	West Virginia	80.1
51	Kentucky	91.4

Survival 

Percent of cases still alive five years after diagnosis

Top 5		
1	Connecticut	27.1%
2	Minnesota	26.9%
3	New Jersey	25.9%
4	Rhode Island	25.9%
5	New York	25.7%
Bottom 5		
42	Alaska	18.4%
42	Kentucky	18.4%
43	West Virginia	18.3%
44	Louisiana	18.1%
45	Mississippi	17.6%
46	Alabama	17.1%

Early Diagnosis 

Percent of cases diagnosed at an early stage

Top 5		
1	Massachusetts	28.8%
2	Wyoming	28.6%
3	Connecticut	27.2%
4	New York	26.1%
5	Delaware	26.0%
Bottom 5		
45	Kentucky	19.8%
46	Oklahoma	19.8%
47	New Mexico	19.0%
48	Alaska	18.6%
49	Hawaii	18.0%

Surgery 


Percent of cases undergoing surgery as part of their first course of treatment

Top 5		
1	Massachusetts	30.7%
2	Rhode Island	28.2%
3	New York	27.7%
4	Connecticut	26.5%
5	New Jersey	25.7%
Bottom 5		
45	Alaska	16.0%
46	Arizona	15.9%
47	Wyoming	15.4%
48	Oklahoma	15.2%
49	New Mexico	13.0%

Untreated 


Percent of cases receiving no treatment after being diagnosed

Top 5		
1	North Dakota	7.5%
2	Massachusetts	9.4%
3	Missouri	9.5%
4	Maine	9.6%
5	New Hampshire	10.6%
Bottom 5		
44	Wyoming	22.1%
45	District of Columbia	22.1%
46	California	22.9%
47	New Mexico	23.8%
48	Arizona	29.5%

Screening 

Percent of those at high risk receiving screening

Top 5		
1	Massachusetts	18.5%
2	Vermont	13.8%
3	New Hampshire	12.1%
4	Kentucky	11.7%
5	Wisconsin	10.8%
Bottom 5		
45	Texas	2.2%
46	Oklahoma	1.7%
47	New Mexico	1.5%
48	California	1.2%
49	Nevada	1.0%

Smoking 

Percent of adults who are current smokers

Top 5		
1	Utah	9.0%
2	California	11.2%
3	Washington	12.0%
4	Connecticut	12.2%
5	Maryland	12.6%
Bottom 5		
47	Tennessee	20.7%
48	Indiana	21.1%
49	Arkansas	22.7%
50	Kentucky	23.4%
51	West Virginia	25.3%