




Tobacco Use in Populations with Mental Illness and/or Substance Use Disorders

**Pat McKone, Regional Senior Director
American Lung Association of the Upper Midwest
Tobacco Control Programs and Policy**


Current Smoking among Adults Aged 18 or Older, by Past Month Serious Psychological Distress Status: NHIS, 1997 to 2011



* Difference between estimate and estimate for 2011 is statistically significant at the .05 level.

Prevalence of Tobacco Use by Mental Health or Substance Use Disorder

Three Fourths of Smokers have a Past or Present Problem with



or Addiction

Mental Illness


Lasser et al., 2000; Data from National Comorbidity Study

Sources: NCS; Lasser, 2000; NESARC; Grant, 2004; NSDUH; MMWR, 2013

Smoking and binge drinking among adults in Minnesota: the current trend




Percent of adults who are current smokers, by binge drinking status,
Minnesota, 2011-2014




Source: Minnesota Behavior Risk Factor Surveillance System, 2011-2014

Mental Health and Tobacco Use among Minnesota Youth

Percent who used tobacco in past 30 days, by presence or absence of mental health indicator:
Minnesota, Grade 11, 2013



Percent who used tobacco in past 30 days, by presence or absence of substance use in past 30 days:
Minnesota, Grade 11, 2013



Tobacco use and substance use among Minnesota Youth





**Pat McKone, Regional Senior Director
American Lung Association of the Upper Midwest
Tobacco Control Programs and Policy**

Pat.McKone@lung.org